



2

NAVAL POSTGRADUATE SCHOOL

Monterey, California

DTIC
ELECTE
MAR 03 1994
S F D



A COMPARISON OF PREDICTORS FOR FIRST-GUESS WIND SPEED ERRORS

Donald P. Gaver
Patricia A. Jacobs

December 1993

Approved for public release; distribution is unlimited.

Prepared for:
Naval Research Laboratory-West
Monterey, CA 93943-5006

3693

94-06907



94 2 2 001

NAVAL POSTGRADUATE SCHOOL
MONTEREY, CA 93943-5000


Rear Admiral T. A. Mercer
Superintendent


Harrison Shull
Provost

This report was prepared for and funded by Naval Research Laboratory-West.

Reproduction of all or part of this report is authorized.


This report was prepared by:



DONALD P. GAVAR, JR.
Professor of Operations Research


PATRICIA A. JACOBS
Professor of Operations Research

Reviewed by:

Released by:


PETER PURDUE
Professor and Chairman
Department of Operations Research


PAUL J. MARTO
Dean of Research

Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

DTIC QUALITY INSPECTED 2

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE December 1993	3. REPORT TYPE AND DATES COVERED Technical Report		
4. TITLE AND SUBTITLE A Comparison of predictors for first-guess wind speed errors		5. FUNDING NUMBERS RH2G5		
6. AUTHOR(S) Donald P. Gaver and Patricia A. Jacobs				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Postgraduate School Monterey, CA 93943		8. PERFORMING ORGANIZATION REPORT NUMBER NPS-OR-93-020		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Naval Research Laboratory-West Monterey, CA 93943-5006		10. SPONSORING / MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) Numerical meteorological models are used to assist in the prediction of weather. Each run of a numerical model produces forecasts of meteorological variables which are used as preliminary predictions of the future values of these variables. These initial predictions are referred to as <i>first-guess</i> values. Estimation of the mean-square first-guess error is required in the optimal interpolation process in the numerical prediction of atmospheric variables. Several predictors for the mean-square error of the first-guess wind speeds are studied. The results suggest that prediction using observed covariates tend to be better than those using first-guess covariates. However, observed covariates are not always available. Predictions using first-guess covariates are better at the 250 mb level than the 850 or 500 mb levels. Of those first-guess covariates studied, first-guess wind speed appears to be the best.				
14. SUBJECT TERMS Gaussian model with log-linear scale parameter; nonparametric models; prediction of mean square errors; first-guess errors in meteorological models; generalized linear regression			15. NUMBER OF PAGES 36	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	

A Comparison of Predictors For First-Guess Wind Speed Errors

by P. A. Jacobs and D. P. Gaver

Abstract

Numerical meteorological models are used to assist in the prediction of weather. Each run of a numerical model produces forecasts of meteorological variables which are used as preliminary predictions of the future values of these variables. These initial predictions are referred to as *first-guess* values. Estimation of the mean-square first-guess error is required in the optimal interpolation process in the numerical prediction of atmospheric variables. Several predictors for the mean-square error of the first-guess wind speeds are studied. The results suggest that prediction using observed covariates tend to be better than those using first-guess covariates. However, observed covariates are not always available. Predictions using first-guess covariates are better at the 250 mb level than the 850 or 500 mb levels. Of those first-guess covariates studied, first-guess wind speed appears to be the best.

1. INTRODUCTION AND SUMMARY

Numerical meteorological models are used to assist in the prediction of weather. Each run of a numerical model produces forecasts of meteorological variables which are used as preliminary predictions of future values of these variables. These initial predictions are referred to as first-guess values. In this paper first-guess values will refer to the most recent 12-hour forecasts.

In certain areas of the world, observations of forecasted variables become available. Prior to the next run of the numerical model a multivariate optimal interpolation analysis updates a first-guess value of a variable by

adding to it a weighted observed value of the variable if it is available. The weight multiplying the observed value depends on estimates of the mean-squared error of the first-guess value and the mean-squared error of the observation; cf. Goerss et al., [1991, a, b]. Thus it is of importance to predict the first-guess mean-squared errors.

The general problem of modeling and predicting mean-square errors is important but not widely studied; see Davidian and Carroll (1987), Nelder and Lee (1992), Aitken (1987), McCullagh and Nelder (1983).

In Jacobs and Gaver (1991, 1992) statistical models for the error of the first guess are used to predict mean-square error for first-guess wind components. The models assume that the error of the first guess has a normal distribution with mean 0 and variance which is a function that is log-linear with covariates. Details of the model are presented in Appendix A.

In this paper we use data from February 1991 to compare the predictive ability of various models. The data consist of measurements and 12 hour forecasts (first-guess values) of u and v wind components at the 850 mb, 500 mb and 250 mb pressure levels from 93 stations in North America 25N–75N for the month of February 1991. The forecasts are produced using the NOGAPS Spectral Forecast Model; cf. Hogan et al., (1991). Each station has measurement and first-guess values for every 12 hours; there are some missing observations. These missing values are deleted from the data set. The measurement values are subtracted from the first-guess values to obtain observations of the error of the first-guess value.

Let $U(o;t)$, (respectively $V(o;t)$), be the observed u -wind, (respectively v -wind) component at time t . Let $U(f;t)$, (respectively $V(f;t)$), be the first-guess u -wind (respectively v -wind) component at time t ; $U(f;t)$, (respectively $V(f;t)$)

is the forecasted value of the u -wind (respectively v -wind) component made 12 hours previously. The first-guess error for the u -wind (respectively v -wind) component is

$$Y_U(t) = U(f;t) - U(o;t); \text{ (respectively } Y_V(t) = V(f;t) - V(o;t)). \quad (1.1)$$

The following covariates are considered in the log-linear model for the mean-square error of the first guess.

$$r(o;t) = \left[(U(o;t) - U(o;t-1))^2 + (V(o;t) - V(o;t-1))^2 \right]^{\frac{1}{2}} \quad (1.2)$$

$$w(o;t) = \left[U(o;t)^2 + V(o;t)^2 \right]^{\frac{1}{2}} \quad (1.3)$$

$$r(f;t) = \left[(U(f;t) - U(f;t-1))^2 + (V(f;t) - V(f;t-1))^2 \right]^{\frac{1}{2}} \quad (1.4)$$

$$w(f;t) = \left[U(f;t)^2 + V(f;t)^2 \right]^{\frac{1}{2}}. \quad (1.5)$$

$$r^*(t) = \left[(U(f;t) - U(o;t-1))^2 + (V(f;t) - V(o;t-1))^2 \right]^{\frac{1}{2}} \quad (1.6)$$

$$a(o,U;t) = |U(o;t) - U(o;t-1)|, \quad a(o,V;t) = |V(o;t) - V(o;t-1)| \quad (1.7)$$

$$a(f,U;t) = |U(f;t) - U(f;t-1)|, \quad a(f,V;t) = |V(f;t) - V(f;t-1)| \quad (1.8)$$

$$a^*(U;t) = |U(f;t) - U(o;t-1)|, \quad a^*(V;t) = |V(f;t) - V(o;t-1)| \quad (1.9)$$

$$m(f;t) = \max(U(f;t), V(f;t)) \quad (1.10)$$

The resultant wind $r(o;t)$, (respectively $r(f;t)$ and $r^*(t)$), is a measure of the observed (respectively forecasted), change in the wind. The variable $w(o;t)$, (respectively $w(f;t)$), is the observed, (respectively forecasted), wind speed. Higher wind speeds suggest more activity in the atmosphere. The change in magnitudes $a(o,U;t)$, $a(f,U;t)$ and $a^*(U;t)$ (respectively $a(o,V;t)$, $a(f,V;t)$ and $a^*(V;t)$) will be used to predict $Y_U(t)$, (respectively $Y_V(t)$).

The data are randomly divided into two sets called DA and DB. Maximum likelihood estimates of the parameters of the models using different covariates are computed using data DA (respectively DB). Nonparametric models based on binning are also considered. The models are then used to predict the mean-square first-guess errors in data set DB (respectively DA). Log-likelihood functions and the empirical distribution of the first-guess errors normalized by their predicted mean-square errors are used to evaluate the models' predictive ability. Details are given in Section 2.

In general, models which use observed covariates, e.g. $w(o)$, $a(o)$, have more predictive ability than those that use first-guess covariates, e.g. $w(f)$, $a(f)$, $m(f)$. The models applied at the 250 mb level appear to have more predictive ability than those for 500 mb and 850 mb.

Among the one-variate models for the 250 mb pressure height, the models that statistically appear to have the most predictive ability have as their covariate $w(o)$, $a(o)$ or $r(o)$. Those that have less but some predictive ability have as their covariate a^* , $w(f)$, $m(f)$ or r^* . Finally, one-variate models using variates $r(f)$ and $a(f)$ appear to have little predictive ability. Among those models for the 250 mb pressure height that use one first-guess covariate, $m(f)$ or $w(f)$ appear to have the most predictive ability.

2. THE DATA ANALYSIS

In this section we describe the data analysis. Let $U_i(o;t)$ and $U_i(f;t)$, (respectively $V_i(o;t)$ and $V_i(f;t)$) be the observed and first-guess u -wind (respectively v -wind) component at location $i = 1, \dots, S$ at time t . By data we mean the vector $(U_i(o,t), U_i(f,t), V_i(o,t), V_i(f,t), U_i(o,t-1), U_i(f,t-1), V_i(o,t-1), V_i(f,t-1))$. The data set contains missing values. Vectors containing these missing values are deleted from the data set. Once missing values are deleted, there are 3618 vectors at the 850 mb level, 4100 at the 500 mb level, and 3744 at the 250 mb level. The observed values are subtracted from the first-guess values to obtain observations of the first-guess errors for each wind component

$$Y_i(U;t) = U_i(f;t) - U_i(o;t)$$

$$Y_i(V;t) = V_i(f;t) - V_i(o;t).$$

The remaining data are randomly divided into two sets called DA and DB without regard to the values of the data, the time t , or the location. Thus, data from the same location for different times may be in different data sets. Models are estimated for each pressure level using only covariates for that pressure level. The covariates considered for each wind component appear in Appendix B. The general statistical model is described in Appendix A.

The model is estimated using data sets DA, DB, and all the data for each pressure level. The estimated values for the parameters for selected models appear in Tables 3A, 4A, 3B, 4B, 3C and 4C. Note that the parameter estimates are usually positive. Hence increased values of the covariates are associated with higher variance of the first-guess errors.

The models estimated from DA (respectively DB) are used to predict the first-guess errors in data set DB (respectively DA). One measure used for

assessing a model's goodness of fit and predictive ability is the value of $\bar{\ell}$, the log-likelihood function up to addition of constants given in Appendix A (A.4); the log-likelihood for predicting mean-square errors in DB using a model estimated using DA uses the first-guess error and covariate(s) from DB and the parameter estimates from DA. Values of $\bar{\ell}$ are computed for data DA (respectively DB) using the parameters estimated using DB (respectively DA); these values assess each model's predictive ability. Values of $\bar{\ell}$ are also computed for data DA (respectively DB) using parameters estimated using DA (respectively DB); these values assess each model's goodness of fit.

Tables 1A, 1B, 1C present the values of $\bar{\ell}$ for one-variate models for the different pressure levels. Also displayed are the values of $\bar{\ell}$ for a model in which the first-guess errors are independent normally distributed with mean 0 and constant variance e^{α} .

Tables 2A, 2B, 2C present values for $\bar{\ell}$ for two-variate models.

Compare the value of $\bar{\ell}_c$ for the model with constant variance (no covariates) for DA (respectively DB) fit using DA (respectively DB) with the values of $\bar{\ell}$ for DA (respectively DB) using models with parameters estimated using the other half of the data DB (respectively DA). A value of $\bar{\ell}$ greater than $\bar{\ell}_c$ indicates that the corresponding model fit with the other half of the data describes the data better than the best constant variance model fit with the same data it is used to summarize. For 850 mb data those one-variate models for which $\bar{\ell} > \bar{\ell}_c$ for DA and DB for both wind components are those with variate r^* , a^* , r , and $w(o)$. For 500 mb the one-variate models are those with variate r^* , $a(o)$, r , $w(o)$, and $w(f)$. For 250 mb, the models are those with variate r^* , $a(o)$, r , $w(o)$, $w(f)$, and $m(f)$.

To compare the predictive ability of the models, the fraction of increase in $\bar{\ell}$, $(\bar{\ell} - \bar{\ell}_c)/|\bar{\ell}_c|$ is computed where $\bar{\ell}_c$ is the maximum value of $\bar{\ell}$ for the constant variance model (with no covariates) estimated using data DA (respectively DB) compared to the value of $\bar{\ell}$ for DA using one-variate models estimated using the other half of the data DB (respectively DA). The values of percentage of increase appear in Table 5 for the one-variable models with variate r^* , a^* , r , $w(o)$, $w(f)$, and $m(f)$. Note that the fraction increase tends to be larger for the 250 mb for the covariates using observed data, r , $w(o)$, and $a(o)$. The fraction also tends to be larger for the first-guess covariates $w(f)$ and $m(f)$ at the 250 mb level.

Another measure of predictability is the distribution of the first-guess errors divided by their predicted standard deviations. Table 6 displays the moments of the first-guess errors of the wind components DA (respectively DB) divided by the standard deviations that are predicted for it using the model fit using data of DB (respectively DA). Recall that the models assume that these errors are normally distributed with mean 0. Thus, if a model were perfect then the mean (respectively standard deviation, skewness, and kurtosis) of the normalized first-guess errors would be 0, (respectively 1, 0 and 3). Of particular interest is the kurtosis. In this application, the kurtosis can be thought of as a measure of the variability of the variance (cf. Cramér page 356). Hence, the smaller the kurtosis, the better the prediction of the model.

Table 6 presents not only results for the model of Appendix A with various covariates but also results for a nonparametric one-variate model. This nonparametric model is as follows. The data in DA (respectively DB) are binned into N bins according to the value of the ordered covariate. For each bin, the mean of the square of the wind speed errors corresponding to the

covariates in that bin is computed. To evaluate the predictive ability of the model, the other data set DB (respectively DA) is used. The predicted mean-square error for a data point in DB (respectively DA) is the mean of the square of the wind speed errors for the bin determined from DA (respectively DB) the data point's covariate lies in.

Table 6 presents selected results using data for the 250 mb level. Results for parametric models of Appendix A with parameters estimated by maximum likelihood (MLE) and the nonparametric models with bins are presented. Also displayed are the sample moments of the first-guess errors in the row labeled "none". Displayed in the row labeled "constant" are the sample moments for the first-guess errors divided by the predicted standard deviation for a model with constant variance e^α fit using the other half of the data.

The values of the kurtosis suggest the following. Once again, models using the observed covariates $a(o)$ and $r(o)$ appear to make the best predictors. Among the first-guess covariates, $m(f)$ and $w(f)$ appear to have comparable predictive ability. If a nonparametric model using a first-guess variate is being considered, then using first-guess wind speed as the covariate seems to be a good choice.

REFERENCES

- Aitken, M., (1987). "Modeling variance heterogeneity in normal regression using GLIM." *Appl Stat.*, 36, pp. 332-339.
- Cox, D. R. and D. V. Hinkley, *Theoretical Statistics*, Chapman and Hall, London, 1974.
- Cramér, H., *Mathematical Methods of Statistics*, Princeton University Press, Princeton, NJ, 1946.
- Davidian, M., and R. J. Carroll (1987). "Variance function estimation," *J. Am Statist. Ass.*, 82, pp. 1079-1091.
- de Bruijn, N. G., *Asymptotic Methods in Analysis*, Interscience, New York, 1958.
- Easton, G. S., "Location compromise maximum likelihood estimators," in *Configural Polysampling*, ed. S. Morgenthaler and J. W. Tukey, Wiley, New York, 1991, pp. 157-192.
- Goerss, J. S. and P. A. Phoebus (1991a). *The Multivariate Optimum Interpolation Analysis of Meteorological Data at FNOC*. NOARL Report Number 31, Naval Oceanographic and Atmospheric Research Laboratory, Stennis Space Center, MS.
- Goerss, J. S. and P. A. Phoebus (1991b). "The Navy's operational atmospheric analysis." to appear in *Weather and Forecasting*.
- Hogan, T. F., and T. E. Rosmond (1991). "The description of the Navy operational global atmospheric prediction system's spectral forecast model." *Monthly Weather Review*, 119, No. 8., pp. 1786-1815.
- Jacobs, P. A. and D. P. Gaver, "Preliminary results from the analysis of wind component error." *Naval Postgraduate School Technical Report NPSOR-91-029*, September, 1991.
- Jacobs, P. A. and D. P. Gaver, "Preliminary results from the analysis of wind component error—July Data." *Naval Postgraduate School Technical Report*, to appear.
- McCullagh and J. A. Nelder, *Generalized Linear Regression*, Chapman and Hall, New York, 1983.
- Nelder, J. A. and Y. Lee (1992), "Likelihood, quasi-likelihood and pseudo-likelihood: some comparisons." *J. R. Statist. Soc., B*, 54, No. 1, pp. 273-284.

APPENDIX A

THE STATISTICAL MODEL

In this Appendix we describe the statistical model. Let $U_i(o;t)$ (respectively $V_i(f;t)$) denote the observed u -wind component (respectively first-guess wind component) at location i at time t ; $i = 1, \dots, S$. Let $V_i(o;t)$, (respectively $V_i(f;t)$) denote the observed v -wind component (respectively first-guess wind component) at location i at time t . The first-guess error of the u -wind (respectively v -wind) component at location i at time t is

$$Y_i(U;t) = U_i(o;t) - U_i(f;t)$$

(respectively,

$$Y_i(V;t) = V_i(o;t) - V_i(f;t).$$

(A.1)

The model is that $\{Y_i(U;t), i = 1, \dots, S\}$ and $\{Y_i(V;t), i = 1, \dots, S\}$ are independent random variables having a normal distribution with mean 0. The variance of $Y_i(U;t)$ is log-linear with a number of covariates. That is

$$\begin{aligned} & \text{Var}[Y_i(U;t) | X_i(1;t) = x_i(1), \dots, X_i(p;t) = x_i(p)] \\ &= \exp \left\{ \alpha + \sum_{j=1}^p \beta_j(t) x_i(j;t) \right\}. \end{aligned} \quad (\text{A.2})$$

The likelihood function for this model is (up to multiplication by constants)

$$\begin{aligned} & L(\alpha, \beta_1, \dots, \beta_p) \\ &= \prod_t \prod_i \exp \left\{ -\frac{1}{2} \left[\alpha + \sum_{j=1}^p \beta_j(t) x_i(j;t) \right] \right\} \exp \left\{ -\frac{1}{2} y^2 \exp \left\{ -\left(\alpha + \sum_{j=1}^p \beta_j(t) x_i(j;t) \right) \right\} \right\} \end{aligned} \quad (\text{A.3})$$

The log-likelihood function is (up to addition by constants)

$$\begin{aligned} & \tilde{l}(\alpha, \beta_1, \dots, \beta_p) \\ &= \sum_t \sum_i -\frac{1}{2} \left[\alpha + \sum_{j=1}^p \beta_j(t) x_i(j;t) \right] - \frac{1}{2} y^2 \exp \left\{ - \left(\alpha + \sum_{j=1}^p \beta_j(t) x_i(j;t) \right) \right\} \end{aligned} \quad (\text{A.4})$$

The recursive procedure used to estimate the parameters $(\alpha, \beta_1, \dots, \beta_p)$ is described in Gaver and Jacobs [1991].

APPENDIX B

THE COVARIATES

In this Appendix we list the covariates that were considered. As before let $U_i(o;t)$ and $U_i(f;t)$, (respectively $V_i(o;t)$ and $V_i(f;t)$) denote the observed and first-guess u -wind (respectively v -wind) component at time t for location i , $i = 1, \dots, S$. The covariates considered for the first-guess error of the u -wind component are

$$a_i(o, U; t) = |U_i(o; t) - U_i(o; t-1)|$$

$$a_i(f, U; t) = |U_i(f; t) - U_i(f; t-1)|$$

$$a_i^*(U; t) = |U_i(f; t) - U_i(o; t-1)|$$

$$w_i(o; t) = [U_i(o; t)^2 + V_i(o; t)^2]$$

$$w_i(f; t) = [U_i(f; t)^2 + V_i(f; t)^2]$$

$$r_i(o; t) = \left[[U_i(o; t) - U_i(o; t-1)]^2 + [V_i(o; t) - V_i(f; t-1)]^2 \right]^{\frac{1}{2}}$$

$$r_i(f; t) = \left[[U_i(f; t) - U_i(f; t-1)]^2 + [V_i(o; t) - V_i(f; t-1)]^2 \right]^{\frac{1}{2}}$$

$$r_i^*(t) = \left[[U_i(f; t) - U_i(o; t-1)]^2 + [V_i(f; t) - V_i(o; t-1)]^2 \right]^{\frac{1}{2}}$$

$$m(f; t) = \max(U(f; t), V(f; t)).$$

The covariates considered for the first-guess error of the v -wind component are

$$a_i(o, V; t) = |V_i(o; t) - V_i(o; t-1)|$$

$$a_i(f, V; t) = |V_i(f; t) - V_i(f; t-1)|$$

$$a_i^*(V; t) = |V_i(f; t) - V_i(o; t-1)|$$

$w_i(o; t), w_i(f; t), r_i(o; t), r_i(f; t),$ and $r_i^*(t).$

TABLE 1A
Log-Likelihood
850 mb. Height/February Data

One-Variate Models

Wind Comp	Data	Model	Const	$r^*(t)$	$r(f;t)$	$a(o;t)$	$a(f;t)$	$a^*(t)$	$r(t)$	$w(o;t)$	$w(f;t)$	$m(f;t)$
u	A	A	-6226.6	-6183.6	-6226.6	-6109.6	-6226.6	-6167.1	-6162.3	-6174.4	-6217.1	-6221.5
	B	B	-6439.8	-6331.3	-6434.9	-6327.7	-6403.8	-6307.6	-6377.8	-6312.1	-6412.9	-6439.0
	B	A	-6452.9	-6347.4	-6452.5	-6341.0	-6450.1	-6319.6	-6390.2	-6333.9	-6426.7	-6455.0
	A	B	-6238.8	-6196.0	-6241.6	-6121.9	-6257.0	-6176.1	-6173.8	-6193.8	-6228.6	-6235.7
v	A	A	-6445.9	-6356.9	-6441.4	-6440.5	-6443.6	-6362.4	-6406.9	-6275.3	-6435.8	-6442.9
	B	B	-6343.7	-6297.4	-6343.1	-6287.4	-6343.5	-6317.6	-6258.1	-6201.8	-6327.9	-6343.7
	B	A	-6346.5	-6302.2	-6347.3	-6316.4	-6349.4	-6326.4	-6271.6	-6207.2	-6332.2	-6336.7
	A	B	-6448.8	-6362.4	-6446.0	-6470.7	-6450.5	-6373.7	-6422.0	-6280.7	-6440.4	-6451.4

$$r(f;t) = \left[(U(f;t) - U(f;t-1))^2 + (V(f;t) - V(f;t-1))^2 \right]^{1/2}$$

$$a(f;t) = |U(f;t) - U(f;t-1)|$$

$$r^*(t) = \left[(U(o;t-1))^2 + (V(o;t-1))^2 \right]^{1/2}$$

$$a(o;t) = |U(o;t) - U(o;t-1)| \quad \text{for } u - \text{wind component error}$$

$$a(o;t) = |V(o;t) - V(o;t-1)| \quad \text{for } v - \text{wind component error}$$

$$a^*(t) = |U(f;t) - U(o;t-1)| \quad \text{for } u - \text{wind component error}$$

$$a^*(t) = |V(f;t) - V(o;t-1)| \quad \text{for } v - \text{wind component error}$$

$$r(t) = \left[(U(o;t) - U(o;t-1))^2 + (V(o;t) - V(o;t-1))^2 \right]^{1/2}$$

$$w(o;t) = \left[U(o;t)^2 + V(o;t)^2 \right]^{1/2}$$

$$w(f;t) = \left[U(f;t)^2 + V(f;t)^2 \right]^{1/2}$$

$$m(f;t) = \max(U(f;t), V(f;t))$$

TABLE 2A
Log-Likelihood
850 mb. Height/February Data

Two-Variate Models

Wind Comp	Data	Model	Const	$r^*(t), a(o;t)$	$r(f;t), a(f;t)$	$r^*(t), a^*(t)$	$r^*(t), a(f;t)$	$r(t), w(o;t)$	$r(t), w(f;t)$
u	A	A	-6226.6	-6085.7	-6226.6	-6164.6	-6177.0	-6137.0	-6160.4
	B	B	-6439.8	-6240.0	-6394.5	-6299.0	-6329.4	-6294.8	-6362.8
	B	A	-6452.9	-6262.5	-6449.6	-6312.4	-6367.9	-6318.9	-6379.3
	A	B	-6238.8	-6102.3	-6263.4	-6174.8	-6202.6	-6158.7	-6173.6
v	A	A	-6445.9	-6356.8	-6441.1	-6353.6	-6344.2	-6260.1	-6404.2
	B	B	-6343.7	-6271.4	-6339.1	-6297.1	-6267.8	-6181.7	-6257.1
	B	A	-6346.5	-6304.1	-6345.6	-6306.5	-6277.0	-6190.3	-6270.8
	A	B	-6448.8	-6396.3	-6448.7	-6365.0	-6355.1	-6270.2	-6419.7

$$r(f;t) = \left[(U(f;t) - U(f;t-1))^2 + (V(f;t) - V(f;t-1))^2 \right]^{1/2}$$

$$a(f;t) = |U(f;t) - U(f;t-1)|$$

$$r^*(t) = \left[(U(f;t) - U(o;t-1))^2 + (V(f;t) - V(o;t-1))^2 \right]$$

$$a(o;t) = |U(o;t) - U(o;t-1)| \quad \text{for } u - \text{wind component error}$$

$$a(o;t) = |V(o;t) - V(o;t-1)| \quad \text{for } v - \text{wind component error}$$

$$a^*(t) = |U(f;t) - U(o;t-1)| \quad \text{for } u - \text{wind component error}$$

$$a^*(t) = |V(f;t) - V(o;t-1)| \quad \text{for } v - \text{wind component error}$$

$$r(t) = \left[(U(o;t) - U(o;t-1))^2 + (V(o;t) - V(o;t-1))^2 \right]^{1/2}$$

$$w(o;t) = \left[U(o;t)^2 + V(o;t)^2 \right]^{1/2}$$

$$w(f;t) = \left[U(f;t)^2 + V(f;t)^2 \right]^{1/2}$$

TABLE 3A
One-Variate Models
Parameter Estimates
(Standard Errors)
850 mb. February Data

Wind Comp	Data Set	$r^*(t)$		$r(f;t)$		$a(o;t)$		$a(f;t)$		$a^*(t)$	
		α	β	α	β	α	β	α	β	α	β
u	A	2.08 (0.06)	0.04 (0.007)	2.44 (0.06)	0.0005 (0.007)	2.00 (0.05)	0.08 (0.008)	2.44 (0.05)	0.002 (0.01)	2.14 (0.05)	0.07 (0.009)
	B	2.00 (0.06)	0.07 (0.007)	2.46 (0.06)	0.02 (0.007)	2.15 (0.05)	0.08 (0.008)	2.33 (0.05)	0.06 (0.01)	2.10 (0.05)	0.09 (0.009)
	All	2.03 (0.04)	0.06 (0.005)	2.45 (0.04)	0.009 (0.005)	2.08 (0.04)	0.08 (0.006)	2.37 (0.03)	0.04 (0.008)	2.11 (0.04)	0.08 (0.006)
v	A	2.07 (0.06)	0.06 (0.007)	2.47 (0.05)	0.02 (0.007)	2.48 (0.05)	0.02 (0.008)	2.51 (0.05)	0.01 (0.008)	2.20 (0.05)	0.06 (0.007)
	B	2.13 (0.06)	0.05 (0.007)	2.47 (0.06)	0.006 (0.008)	2.22 (0.05)	0.06 (0.008)	2.52 (0.05)	-0.004 (0.008)	2.29 (0.05)	0.04 (0.008)
	All	2.10 (0.04)	0.05 (0.005)	2.47 (0.04)	0.01 (0.005)	2.35 (0.03)	0.04 (0.005)	2.51 (0.03)	0.005 (0.006)	2.24 (0.04)	0.05 (0.005)

$$r(f;t) = \left[(U(f;t) - U(f;t-1))^2 + (V(f;t) - V(f;t-1))^2 \right]^{1/2}$$

$$a(f;t) = |U(f;t) - U(f;t-1)|$$

$$r^*(t) = \left[(U(o;t) - U(o;t-1))^2 + (V(o;t) - V(o;t-1))^2 \right]$$

$$a(o;t) = |U(o;t) - U(o;t-1)|, \quad a^*(t) = |U(f;t) - U(o;t-1)|$$

TABLE 4A
Two-Variate Models
Parameter Estimates
(Standard Errors)
850 mb. February Data

Wind Comp	Data Set	$\log MSE = \alpha + \beta_1 r^*(t) + \beta_2 a(o;t)$			$\log MSE = \alpha + \beta_1 r_f(t) + \beta_2 a(f;t)$			$\log MSE = \alpha + \beta_1 r^*(t) + \beta_2 a^*(t)$		
		α	β_1	β_2	α	β_1	β_2	α	β_1	β_2
u	A	1.75 (0.07)	0.04 (0.007)	0.08 (0.008)	2.13 (0.06)	0.05 (0.008)	-0.03 (0.012)	2.08 (0.06)	0.02 (0.01)	0.05 (0.01)
	B	1.65 (0.07)	0.06 (0.007)	0.07 (0.008)	1.98 (0.06)	0.06 (0.008)	0.02 (0.01)	1.99 (0.06)	0.03 (0.009)	0.07 (0.01)
	All	1.70 (0.05)	0.05 (0.005)	0.08 (0.006)	2.04 (0.05)	0.06 (0.006)	-0.004 (0.008)	2.03 (0.04)	0.02 (0.007)	0.06 (0.009)
v	A	2.08 (0.07)	0.06 (0.007)	-0.002 (0.009)	2.47 (0.06)	0.02 (0.01)	-0.009 (0.02)	2.09 (0.06)	0.04 (0.01)	0.03 (0.01)
	B	2.04 (0.07)	0.03 (0.008)	0.04 (0.009)	2.45 (0.06)	0.03 (0.02)	-0.03 (0.02)	2.13 (0.06)	0.05 (0.01)	-0.007 (0.01)
	All	2.05 (0.05)	0.05 (0.005)	0.02 (0.006)	2.46 (0.04)	0.03 (0.01)	-0.02 (0.01)	2.10 (0.04)	0.05 (0.009)	0.01 (0.01)

$$r(f;t) = \left[(u(f;t) - u(f;t-1))^2 + (v(f;t) - v(f;t-1))^2 \right]^{1/2}$$

$$a(f;t) = |u(f;t) - u(f;t-1)|$$

$$r^*(t) = \left[(u(f;t) - u(o;t-1))^2 + (v(f;t) - v(o;t-1))^2 \right]$$

$$a(o;t) = |u(o;t) - u(o;t-1)|, \quad a^*(t) = |u(f;t) - u(o;t-1)|$$

TABLE 1B
Log-Likelihood
500 mb. Height/February Data

One-Variate Models

Wind Comp	Data	Model	Const	$r^*(t)$	$r(f;t)$	$a(o;t)$	$a(f;t)$	$a^*(t)$	$r(t)$	$w(o;t)$	$w(f;t)$	$m(f;t)$
u	A	A	-7894.4	-7865.0	-7877.6	-7699.3	-7876.1	-7866.0	-7732.5	-7846.3	-7866.9	-7860.5
	B	B	-7965.1	-7816.5	-7928.9	-7711.9	-7963.0	-7854.9	-7780.2	-7890.7	-7936.8	-7923.2
	B	A	-7966.4	-7854.2	-7933.2	-7714.0	-7973.1	-7871.6	-7780.7	-7893.5	-7937.9	-7925.4
	A	B	-7895.6	-7897.9	-7881.8	-7701.0	-7885.5	-7878.3	-7732.9	-7848.9	-7868.0	-7862.9
v	A	A	-7720.5	-7702.4	-7716.8	-7624.9	-7709.3	-7695.9	-7616.2	-7683.2	-7696.0	-7706.7
	B	B	-7849.7	-7841.9	-7847.6	-7723.5	-7845.6	-7842.2	-7740.4	-7780.6	-7827.2	-7828.2
	B	A	-7853.8	-7849.2	-7852.2	-7727.3	-7853.3	-7854.3	-7744.2	-7787.3	-7830.9	-7832.9
	A	B	-7724.5	-7709.1	-7721.2	-7628.3	-7715.9	-7706.3	-7619.8	-7689.3	-7699.5	-7711.2

$$r(f;t) = \left[(U(f;t) - U(f;t-1))^2 + (V(f;t) - V(f;t-1))^2 \right]^{1/2}$$

$$a(f;t) = |U(f;t) - U(f;t-1)|$$

$$r^*(t) = \left[(U(f;t) - U(o;t-1))^2 + (V(f;t) - V(o;t-1))^2 \right]$$

$$a(o;t) = |U(o;t) - U(o;t-1)| \quad \text{for } u - \text{wind component error}$$

$$a(o;t) = |V(o;t) - V(o;t-1)| \quad \text{for } v - \text{wind component error}$$

$$a^*(t) = |U(f;t) - U(o;t-1)| \quad \text{for } u - \text{wind component error}$$

$$a^*(t) = |V(f;t) - V(o;t-1)| \quad \text{for } v - \text{wind component error}$$

$$r(t) = \left[(U(o;t) - U(o;t-1))^2 + (V(o;t) - V(o;t-1))^2 \right]^{1/2}$$

$$w(o;t) = \left[U(o;t)^2 + V(o;t)^2 \right]^{1/2}$$

$$w(f;t) = \left[U(f;t)^2 + V(f;t)^2 \right]^{1/2}$$

$$m(f;t) = \max(U(f;t), V(f;t))$$

TABLE 2B
Log-Likelihood
500 mb. Height
February Data

Two-Variate Models

Wind Comp	Data	Model	Const	$\hat{r}(t), a(o;t)$	$r(f;t), a(f;t)$	$\hat{r}^*(t), \hat{a}^*(t)$	$\hat{r}^*(t), a(f;t)$	$r(t), w(o;t)$	$r(t), w(f;t)$
u	A	A	-7894.4	-7693.4	-7873.2	-7860.2	-7860.3	-7718.3	-7724.1
	B	B	-7965.1	-7676.8	-7923.7	-7807.4	-7813.6	-7762.7	-7772.2
	B	A	-7966.4	-7689.2	-7950.9	-7843.4	-7873.3	-7763.1	-7772.6
	A	B	-7895.6	-7704.0	-7893.6	-7891.5	-7905.7	-7718.7	-7724.5
v	A	A	-7720.5	-7622.3	-7706.6	-7695.7	-7701.5	-7601.9	-7609.0
	B	B	-7849.7	-7722.0	-7845.0	-7841.3	-7841.9	-7711.0	-7730.8
	B	A	-7853.8	-7733.8	-7853.9	-7853.6	-7850.4	-7716.5	-7734.6
	A	B	-7724.5	-7633.9	-7714.1	-7706.2	-7708.9	-7607.0	-7612.6

$$r(f;t) = \left[(U(f;t) - U(f;t-1))^2 + (V(f;t) - V(f;t-1))^2 \right]^{1/2}$$

$$a(f;t) = |U(f;t) - U(f;t-1)|$$

$$\hat{r}^*(t) = \left[(U(f;t) - U(o;t-1))^2 + (V(f;t) - V(o;t-1))^2 \right]$$

$$a(o;t) = |U(o;t) - U(o;t-1)| \quad \text{for } u - \text{wind component error}$$

$$a(o;t) = |V(o;t) - V(o;t-1)| \quad \text{for } v - \text{wind component error}$$

$$\hat{a}^*(t) = |U(f;t) - U(o;t-1)| \quad \text{for } u - \text{wind component error}$$

$$\hat{a}^*(t) = |V(f;t) - V(o;t-1)| \quad \text{for } v - \text{wind component error}$$

$$r(t) = \left[(U(o;t) - U(o;t-1))^2 + (V(o;t) - V(o;t-1))^2 \right]^{1/2}$$

$$w(o;t) = \left[U(o;t)^2 + V(o;t)^2 \right]^{1/2}$$

$$w(f;t) = \left[U(f;t)^2 + V(f;t)^2 \right]^{1/2}$$

TABLE 3B
One-Variate Models
Parameter Estimates
(Standard Errors)
500 mb. February Data

Wind Comp	Data Set	$r^*(t)$		$r(f;t)$		$a(o;t)$		$a(f;t)$		$a^*(t)$		$m(f;t)$	
		α	β	α	β	α	β	α	β	α	β	α	β
u	A	2.56 (0.06)	0.03 (0.005)	2.63 (0.06)	0.02 (0.006)	2.31 (0.05)	0.07 (0.006)	2.68 (0.05)	0.03 (0.008)	2.64 (0.05)	0.03 (0.007)	2.55 (0.06)	0.02 (0.003)
	B	2.25 (0.06)	0.06 (0.005)	2.60 (0.06)	0.02 (0.006)	2.28 (0.05)	0.08 (0.006)	2.83 (0.05)	0.01 (0.008)	2.50 (0.05)	0.06 (0.006)	2.53 (0.06)	0.02 (0.003)
	All	2.40 (0.04)	0.04 (0.004)	2.60 (0.04)	0.03 (0.004)	2.29 (0.03)	0.08 (0.004)	2.75 (0.03)	0.02 (0.006)	2.56 (0.03)	0.05 (0.005)	2.54 (0.04)	0.02 (0.002)
v	A	2.54 (0.06)	0.02 (0.005)	2.66 (0.06)	0.01 (0.006)	2.41 (0.05)	0.04 (0.005)	2.63 (0.05)	0.02 (0.007)	2.56 (0.05)	0.03 (0.006)	2.57 (0.06)	0.01 (0.003)
	B	2.69 (0.06)	0.01 (0.005)	2.75 (0.06)	0.008 (0.006)	2.40 (0.05)	0.05 (0.005)	2.75 (0.05)	0.01 (0.006)	2.72 (0.05)	0.01 (0.005)	2.58 (0.06)	0.01 (0.003)
	All	2.62 (0.04)	0.02 (0.004)	2.71 (0.04)	0.01 (0.004)	2.40 (0.03)	0.05 (0.003)	2.69 (0.03)	0.02 (0.004)	2.65 (0.03)	0.02 (0.004)	2.57 (0.04)	0.01 (0.002)

$$r(f;t) = \left[(U(f;t) - U(f;t-1))^2 + (V(f;t) - V(f;t-1))^2 \right]^{1/2}$$

$$a(f;t) = |U(f;t) - U(f;t-1)|$$

$$r^*(t) = \left[(U(f;t) - U(o;t-1))^2 + (V(f;t) - V(o;t-1))^2 \right]$$

$$a(o;t) = |U(o;t) - U(o;t-1)|, \quad a^*(t) = |U(f;t) - U(o;t-1)|$$

TABLE 4B
Two-Variate Models
Parameter Estimates
(Standard Errors)
500 mb. February Data

Wind Comp	Data Set	$\log MSE = \alpha + \beta_1 r^*(t) + \beta_2 a(o;t)$			$\log MSE = \alpha + \beta_1 r_f(t) + \beta_2 a(f;t)$			$\log MSE = \alpha + \beta_1 r^*(t) + \beta_2 a^*(t)$		
		α	β_1	β_2	α	β_1	β_2	α	β_1	β_2
u	A	2.20 (0.07)	0.01 (0.005)	0.07 (0.006)	2.62 (0.06)	0.01 (0.008)	0.02 (0.01)	2.55 (0.06)	0.02 (0.007)	0.02 (0.009)
	B	2.02 (0.06)	0.03 (0.005)	0.07 (0.006)	2.58 (0.06)	0.05 (0.008)	-0.02 (0.01)	2.24 (0.06)	0.04 (0.007)	0.02 (0.008)
	All	2.10 (0.05)	0.02 (0.004)	0.07 (0.004)	2.60 (0.04)	0.03 (0.005)	-0.002 (0.007)	2.39 (0.04)	0.03 (0.005)	-0.02 (0.006)
v	A	2.33 (0.06)	0.009 (0.006)	0.04 (0.005)	2.68 (0.06)	-0.02 (0.01)	0.04 (0.01)	2.55 (0.06)	0.004 (0.008)	0.02 (0.009)
	B	2.45 (0.06)	-0.007 (0.006)	0.05 (0.005)	2.78 (0.06)	-0.009 (0.01)	0.02 (0.01)	2.69 (0.06)	0.008 (0.009)	0.007 (0.009)
	All	2.39 (0.05)	0.001 (0.004)	0.05 (0.004)	2.73 (0.04)	-0.01 (0.008)	0.03 (0.008)	2.62 (0.04)	0.006 (0.006)	0.01 (0.006)

$$r(f;t) = \left[(u(f;t) - u(f;t-1))^2 + (v(f;t) - v(f;t-1))^2 \right]^{1/2}$$

$$a(f;t) = |u(f;t) - u(f;t-1)|$$

$$r^*(t) = \left[(u(f;t) - u(o;t-1))^2 + (v(f;t) - v(o;t-1))^2 \right]^{1/2}$$

$$a(o;t) = |u(o;t) - u(o;t-1)|, \quad a^*(t) = |u(f;t) - u(o;t-1)|$$

TABLE 1C
Log-Likelihood
250 mb. Height/February Data

One-Variate Models

Wind Comp	Data	Model	Const	$r^*(t)$	$r(f;t)$	$a(o;t)$	$a(f;t)$	$\hat{a}^*(t)$	$r(t)$	$w(o;t)$	$w(f;t)$	$m(f;t)$	$U(f;t)$ $V(f;t)$
u	A	A	-9376.4	-9253.8	-9365.4	-8609.8	-9362.6	-9289.9	-8705.7	-8614.6	-9137.6	-8962.3	-9074.1
	B	B	-8957.8	-8897.4	-8927.6	-8559.6	-8950.7	-8946.3	-8637.6	-8600.3	-8890.9	-8795.1	-8911.9
	B	A	-9001.3	-8931.4	-8977.3	-8562.1	-9024.6	-8989.3	-8641.4	-8611.9	-8923.2	-8816.7	-8991.9
	A	B	-9426.8	-9292.8	-9423.7	-8612.3	-9463.7	-9351.3	-8710.3	-8629.4	-9188.4	-8994.9	-9186.6
v	A	A	-8803.2	-8669.4	-8801.6	-8477.6	-8782.9	-8621.7	-8435.0	-8355.6	-8692.4	-8693.1	-8786.2
	B	B	-8916.9	-8835.6	-8864.4	-8566.5	-8833.5	-8829.6	-8565.4	-8507.1	-8886.9	-8877.3	-8913.0
	B	A	-8920.4	-8843.7	-8898.1	-8575.9	-8843.0	-8860.6	-8571.7	-8516.4	-8907.8	-8890.4	-8922.6
	A	B	-8806.6	-8677.3	-8820.1	-8486.7	-8788.3	-8647.8	-8440.6	-8364.2	-8714.4	-8706.5	-8794.8

$$r(f;t) = \left[(U(f;t) - U(f;t-1))^2 + (V(f;t) - V(f;t-1))^2 \right]^{1/2}$$

$$a(f;t) = |U(f;t) - U(f;t-1)|$$

$$\hat{r}^*(t) = \left[(U(f;t) - U(o;t-1))^2 + (V(f;t) - V(o;t-1))^2 \right]$$

$$a(o;t) = |U(o;t) - U(o;t-1)| \quad \text{for } u - \text{wind component error}$$

$$a(o;t) = |V(o;t) - V(o;t-1)| \quad \text{for } v - \text{wind component error}$$

$$\hat{a}^*(t) = |U(f;t) - U(o;t-1)| \quad \text{for } u - \text{wind component error}$$

$$\hat{a}^*(t) = |V(f;t) - V(o;t-1)| \quad \text{for } v - \text{wind component error}$$

$$r(t) = \left[(U(o;t) - U(o;t-1))^2 + (V(o;t) - V(o;t-1))^2 \right]^{1/2}$$

$$w(o;t) = \left[U(o;t)^2 + V(o;t)^2 \right]^{1/2}$$

$$w(f;t) = \left[U(f;t)^2 + V(f;t)^2 \right]^{1/2}$$

$$m(f;t) = \max(U(f;t), V(f;t))$$

TABLE 2C
Log-Likelihood
250 mb. Height
February Data

Two-Variate Models

Wind Comp	Data	Model	Const	$r^*(t), a(o;t)$	$r(f;t), a(f;t)$	$r^*(t), a^*(t)$	$r^*(t), a(f;t)$	$r(t), w(o;t)$	$r(t), w(f;t)$
u	A	A	-9376.4	-8537.9	-9361.5	-9252.4	-9231.0	-8504.0	-8685.4
	B	B	-8957.8	-8553.5	-8900.9	-8896.2	-8864.2	-8502.1	-8581.6
	B	A	-9001.3	-8572.7	-9004.9	-8934.7	-8981.0	-8507.2	-8594.4
	A	B	-9426.8	-8567.0	-9479.4	-9297.6	-9465.5	-8509.3	-8698.6
v	A	A	-8803.2	-8417.8	-8756.1	-8621.6	-8660.9	-8288.9	-8426.5
	B	B	-8916.9	-8565.9	-8828.3	-8821.9	-8818.3	-8415.8	-8541.6
	B	A	-8920.4	-8598.2	-8854.5	-8858.3	-8894.1	-8424.4	-8553.3
	A	B	-8806.6	-8469.7	-8772.4	-8646.8	-8710.8	-8296.5	-8437.9

$$r(f;t) = \left[(U(f;t) - U(f;t-1))^2 + (V(f;t) - V(f;t-1))^2 \right]^{1/2}$$

$$a(f;t) = |U(f;t) - U(f;t-1)|$$

$$r^*(t) = \left[(U(f;t) - U(o;t-1))^2 + (V(f;t) - V(o;t-1))^2 \right]$$

$$a(o;t) = |U(o;t) - U(o;t-1)| \quad \text{for } u - \text{wind component error}$$

$$a(o;t) = |V(o;t) - V(o;t-1)| \quad \text{for } v - \text{wind component error}$$

$$a^*(t) = |U(f;t) - U(o;t-1)| \quad \text{for } u - \text{wind component error}$$

$$a^*(t) = |V(f;t) - V(o;t-1)| \quad \text{for } v - \text{wind component error}$$

$$r(t) = \left[(U(o;t) - U(o;t-1))^2 + (V(o;t) - V(o;t-1))^2 \right]^{1/2}$$

$$w(o;t) = \left[U(o;t)^2 + V(o;t)^2 \right]^{1/2}$$

$$w(f;t) = \left[U(f;t)^2 + V(f;t)^2 \right]^{1/2}$$

TABLE 3C
One-Variate Models
Parameter Estimates
(Standard Errors)
250 mb. February Data

Wind Comp	Data Set	$r^*(t)$		$r(f;t)$		$a(o;t)$		$a(f;t)$		$a^*(t)$		$m(f;t)$	
		α	β	α	β	α	β	α	β	α	β	α	β
u	A	3.50 (0.06)	0.03 (0.003)	3.81 (0.07)	0.02 (0.005)	3.10 (0.05)	0.06 (0.004)	3.88 (0.05)	0.02 (0.005)	3.72 (0.04)	0.03 (0.003)	3.00 (0.06)	0.03 (0.002)
	B	3.34 (0.06)	0.03 (0.004)	3.52 (0.06)	0.02 (0.004)	3.07 (0.05)	0.05 (0.004)	3.88 (0.05)	-0.01 (0.005)	3.66 (0.05)	0.01 (0.004)	3.16 (0.07)	0.02 (0.002)
	All	3.42 (0.04)	0.03 (0.002)	3.67 (0.04)	0.02 (0.003)	3.08 (0.03)	0.05 (0.003)	3.87 (0.03)	0.005 (0.004)	3.68 (0.03)	0.02 (0.003)	3.05 (0.05)	0.03 (0.001)
v	A	3.07 (0.06)	0.04 (0.005)	3.63 (0.06)	0.005 (0.004)	3.13 (0.04)	0.04 (0.003)	3.50 (0.05)	0.02 (0.005)	3.10 (0.05)	0.05 (0.004)	3.14 (0.06)	0.02 (0.002)
	B	3.22 (0.07)	0.03 (0.004)	3.42 (0.06)	0.02 (0.004)	3.10 (0.05)	0.04 (0.002)	3.44 (0.05)	0.03 (0.004)	3.37 (0.05)	0.03 (0.004)	3.34 (0.07)	0.01 (0.002)
	All	3.15 (0.05)	0.03 (0.003)	3.46 (0.03)	0.03 (0.003)	3.12 (0.03)	0.04 (0.002)	3.46 (0.03)	0.03 (0.003)	3.25 (0.04)	0.04 (0.003)	3.23 (0.05)	0.02 (0.001)

$$r(f;t) = \left[(U(f;t) - U(f;t-1))^2 + (V(f;t) - V(f;t-1))^2 \right]^{1/2}$$

$$a(f;t) = |U(f;t) - U(f;t-1)|$$

$$r^*(t) = \left[(U(o;t) - U(o;t-1))^2 + (V(o;t) - V(o;t-1))^2 \right]$$

$$a(o;t) = |U(o;t) - U(o;t-1)|, \quad a^*(t) = |U(f;t) - U(o;t-1)|$$

$$m(f;t) = \max(U(f;t), V(f;t))$$

TABLE 4C
Two-Variate Models
Parameter Estimates
(Standard Errors)
250 mb. February Data

Wind Comp	Data Set	$\log MSE = \alpha + \beta_1 r^*(t) + \beta_2 a(o;t)$			$\log MSE = \alpha + \beta_1 r_f(t) + \beta_2 a(f;t)$			$\log MSE = \alpha + \beta_1 r^*(t) + \beta_2 a^*(t)$		
		α	β_1	β_2	α	β_1	β_2	α	β_1	β_2
u	A	2.73 (0.06)	0.03 (0.003)	0.05 (0.003)	3.83 (0.07)	0.007 (0.006)	0.01 (0.007)	3.50 (0.06)	0.03 (0.004)	0.006 (0.005)
	B	2.93 (0.07)	0.009 (0.004)	0.05 (0.004)	3.61 (0.06)	0.03 (0.004)	-0.03 (0.006)	3.35 (0.06)	0.03 (0.004)	-0.006 (0.005)
	All	2.80 (0.05)	0.02 (0.002)	0.05 (0.003)	3.68 (0.04)	0.02 (0.004)	-0.01 (0.004)	3.42 (0.04)	0.03 (0.003)	0.007 (0.004)
v	A	2.80 (0.06)	0.02 (0.003)	0.03 (0.003)	3.69 (0.07)	-0.04 (0.008)	0.06 (0.008)	3.09 (0.06)	0.001 (0.005)	0.05 (0.006)
	B	3.06 (0.07)	0.003 (0.004)	0.04 (0.003)	3.52 (0.06)	-0.02 (0.007)	0.05 (0.007)	3.25 (0.07)	0.02 (0.006)	0.02 (0.006)
	All	2.90 (0.04)	0.02 (0.002)	0.04 (0.002)	3.58 (0.04)	-0.03 (0.005)	0.05 (0.005)	3.18 (0.04)	0.01 (0.004)	0.03 (0.004)

$$r(f;t) = \left[(u(f;t) - u(f;t-1))^2 + (v(f;t) - v(f;t-1))^2 \right]^{1/2}$$

$$a(f;t) = |u(f;t) - u(f;t-1)|$$

$$r^*(t) = \left[(u(f;t) - u(o;t-1))^2 + (v(f;t) - v(o;t-1))^2 \right]$$

$$a(o;t) = |u(o;t) - u(o;t-1)|, \quad a^*(t) = |u(f;t) - u(o;t-1)|$$

TABLE 5
Percent of Increase
 $(\bar{e} - \bar{e}_c)/|\bar{e}_c|$

One-Variate Models

Pressure Level mb	Wind Comp	Data Set	Model	r	$w(o)$	$a(o)$	r^*	a^*	$w(f)$	$m(f)$
850	u	B	A	0.8	1.6	1.5	1.4	1.9	0.2	-0.2
		A	B	0.8	0.5	1.7	0.5	0.8	-0.03	-0.1
	v	B	A	0.7	2.2	0.4	0.7	0.3	0.2	0.1
		A	B	0.4	2.6	-0.3	1.3	1.1	0.08	-0.09
500	u	B	A	2.3	0.9	3.2	1.4	1.2	0.3	0.03
		A	B	2.0	0.6	2.4	-0.04	0.2	0.3	0.4
	v	B	A	1.3	0.8	1.6	0.006	-0.005	0.2	0.2
		A	B	1.3	0.4	1.2	0.1	0.2	0.3	0.1
250	u	B	A	3.5	3.9	4.4	0.3	-0.4	0.4	3.0
		A	B	7.1	8.0	8.1	0.8	0.3	2.0	4.0
	v	B	A	3.9	4.5	3.8	0.8	0.6	0.1	0.3
		A	B	4.1	5.0	3.6	1.4	1.7	1.0	0.1

TABLE 6
Sample Moments of
First-Guess Wind Speed Errors Divided by Predicted Standard Deviations

Wind Comp	Covariates	Est. Method	Mean	Std.Dev.	Skewness	Kurtosis
u	none	-	0.11	7.04	-1.28	51.1
	constant	MLE	0.02	1.01	-1.24	57.4
	w(f)	MLE	0.02	1.01	-0.73	32.9
	w(f)	2 bins	0.02	1.01	-0.59	34.4
		3 bins	0.02	1.02	-0.98	42.9
	m(f)	MLE	0.01	1.01	1.01	27.6
	m(f)	2 bins	0.02	1.01	-0.63	34.1
		3 bins	0.01	1.03	-1.11	41.0
	a(o)	MLE	0.02	1.00	0.24	6.0
	r(o)	MLE	0.02	1.00	0.17	7.7
	a(f)	MLE	0.02	1.02	-0.92	57.6
	a*	MLE	0.01	1.00	-1.42	64.5
v	none	-	-0.11	6.47	-1.98	35.5
	constant	MLE	-0.02	1.00	-1.98	35.3
	w(f)	MLE	-0.00	1.01	-1.49	28.6
	w(f)	2 bins	-0.00	1.00	-1.28	23.9
		3 bins	-0.01	1.03	-2.04	37.8
	m(f)	MLE	-0.00	1.00	-1.33	24.8
	m(f)	2 bins	-0.02	1.00	-2.00	36.2
		3 bins	-0.01	1.04	-2.14	40.9
	a(o)	MLE	-0.02	1.00	2.08	36.1

$$m(f;t) = \max(U(f;t), V(f;t))$$

$$w(f;t) = [U(f;t)^2 + V(f;t)^2]^{1/2}$$

INITIAL DISTRIBUTION LIST

1. Research Office (Code 08)1
 Naval Postgraduate School
 Monterey, CA 93943-5000

2. Dudley Knox Library (Code 52)2
 Naval Postgraduate School
 Monterey, CA 93943-5002

3. Defense Technical Information Center.....2
 Cameron Station
 Alexandria, VA 22314

4. Department of Operations Research (Code OR).....1
 Naval Postgraduate School
 Monterey, CA 93943-5000

5. Prof. Donald Gaver (Code OR/Gv)5
 Naval Postgraduate School
 Monterey, CA 93943-5000

6. Prof. Patricia Jacobs (Code OR/Jc)5
 Naval Postgraduate School
 Monterey, CA 93943-5000

7. Center for Naval Analyses.....1
 4401 Ford Avenue
 Alexandria, VA 22302-0268

8. Dr. J. Abrahams, Code 1111, Room 6071
 Mathematical Sciences Division, Office of Naval Research
 800 North Quincy Street
 Arlington, VA 22217-5000

9. Dr. David Brillinger.....1
 Statistics Department
 University of California
 Berkeley, CA 94720

10. Prof. Brad Carlin.....1
School of Public Health
University of Minnesota
Mayo Bldg. A460
Minneapolis, MN 55455

11. Prof. H. Chernoff.....1
Department of Statistics
Harvard University
1 Oxford Street
Cambridge, MA 02138

12. Dr. John Copas.....1
Dept. of Mathematics,
University of Birmingham
P. O. Box 363
Birmingham B15 2TT
ENGLAND

13. Professor Sir David Cox.....1
Nuffield College
Oxford, OXI INF
ENGLAND

14. Professor H. G. Daellenbach.....1
Department of Operations Research
University of Canterbury
Christchurch, NEW ZEALAND

15. Dr. S. R. Dalal.....1
Bellcore
445 South Street
Morristown, NJ 07962-1910

16. Dr. D. F. Daley1
Statistic Dept. (I.A.S.)
Australian National University
Canberra, A.C.T. 2606
AUSTRALIA

17. Prof. Bradley Efron.....1
 Statistics Dept.
 Sequoia Hall
 Stanford University
 Stanford, CA 94305

18. Prof. George S. Fishman.....1
 Curr. in OR & Systems Analysis
 University of North Carolina
 Chapel Hill, NC 20742

19. Dr. Neil Gerr.....1
 Office of Naval Research
 Arlington, VA 22217

20. Dr. J. Goerss.....10
 Naval Oceanographic and Atmospheric Laboratory
 Monterey, CA 93943-5006

21. Dr. D. C. Hoaglin.....1
 Department of Statistics
 Harvard University
 1 Oxford Street
 Cambridge, MA 02138

22. Institute for Defense Analysis.....1
 1800 North Beauregard
 Alexandria, VA 22311

23. Prof. J. B. Kadane.....1
 Dept. of Statistics
 Carnegie-Mellon University
 Pittsburgh, PA 15213

24. Dr. Jon Kettenring.....1
 Bellcore
 445 South Street
 Morris Township, NJ 07962-1910

25. Koh Peng Kong.....1
 OA Branch, DSO
 Ministry of Defense
 Blk 29 Middlesex Road
 SINGAPORE 1024

26. Dr. A. J. Lawrence.....1
 Dept. of Mathematics,
 University of Birmingham
 P. O. Box 363
 Birmingham B15 2TT
 ENGLAND

27. Prof. M. Leadbetter.....1
 Department of Statistics
 University of North Carolina
 Chapel Hill, NC 27514

28. Prof. J. Lehoczky.....1
 Department of Statistics
 Carnegie-Mellon University
 Pittsburgh, PA 15213

29. Dr. Colin Mallows.....1
 AT&T Bell Telephone Laboratories
 600 Mountain Avenue
 Murray Hill, NJ 07974

30. Prof. M. Mazumdar.....1
 Dept. of Industrial Engineering
 University of Pittsburgh
 Pittsburgh, PA 15235

31. Prof. Carl N. Morris.....1
 Statistics Department
 Harvard University
 1 Oxford St.
 Cambridge, MA 02138

32. Prof. F. W. Mosteller.....1
 Department of Statistics
 Harvard University
 1 Oxford St.
 Cambridge, MA 02138

33. Operations Research Center, Rm E40-164.....1
 Massachusetts Institute of Technology
 Attn: R. C. Larson and J. F. Shapiro
 Cambridge, MA 02139

34. Prof. Frank Samaniego1
 Statistics Department
 University of California
 Davis, CA 95616

35. Prof. W. R. Schucany.....1
 Dept. of Statistics
 Southern Methodist University
 Dallas, TX 75222

36. Prof. H. Solomon1
 Department of Statistics
 Sequoia Hall
 Stanford University
 Stanford, CA 94305

37. Prof. W. Stuetzle1
 Department of Statistics
 University of Washington
 Seattle, WA 98195

38. Prof. J. R. Thompson.....1
 Dept. of Mathematical Science
 Rice University
 Houston, TX 77001

39. Prof. J. W. Tukey1
 Statistics Dept., Fine Hall
 Princeton University
 Princeton, NJ 08540

40. Dr. D. Vere-Jones.....1
 Dept. of Math, Victoria Univ. of Wellington
 P. O. Box 196
 Wellington
 NEW ZEALAND

41. Prof. David L. Wallace1
 Statistics Dept., University of Chicago
 5734 S. University Ave.
 Chicago, IL 60637

42. Dr. Ed Wegman.....1
George Mason University
Fairfax, VA 22030
43. Dr. P. Welch1
IBM Research Laboratory
Yorktown Heights, NY 10598
44. Prof. Roy Welsch.....1
Sloan School
M.I.T.
Cambridge, MA 02139